

参考文献

- [1] Taoka M, Takashi T, Iwamura Y. Representation of the mid-line trunk, bilateral arms, and shoulders in the monkey post-central somatosensory cortex[J]. Exp Brain Res, 1998,123: 315—322.
- [2] 中华神经学会. 各类脑血管疾病诊断要点[J]. 中华神经科杂志, 1996,29:379—380.
- [3] 杨坚, 乔蕾, 朱琪, 等. 个体化主动康复对脑卒中偏瘫患者运动功能和日常生活活动能力的影响[J]. 中国康复医学杂志, 2007, 22(6):514—517.
- [4] Fugl-Meyer AR, Jääskö L, Leyman I, et al. The post-stroke hemiplegic patient. A method for evaluation of physical performance[J]. Scand J Rehabil Med, 1975,7(1):13—31.
- [5] Platz T, Pinkowski C, van Wijek F, et al. Reliability and validity of arm function assessment with standardized guidelines for the Fugl-Meyer Test, Action Research Arm Test and Box and Block Test: a multicentre study[J]. Clinical Rehabilitation, 2005, 19:404—411.
- [6] Hsueh IP, Lin JH, Jeng JS, et al. Comparison of the psychometric characteristics of the functional independence measure, 5 item Barthel index, and 10 item Barthel index in patients with stroke[J]. Journal of Neurology, Neurosurgery & Psychiatry, 2002,73(2):188—190.
- [7] Collin C, Wade DT, Davies S, Horne V. The Barthel ADL Index: a reliability study[J]. Int Disabil Stud, 1988, 10: 61—63.
- [8] Michaelsen SM, Jacobs S, Roby-Brami A, Levin MF. Compensation for distal impairments of grasping in adults with hemiparesis[J]. Exp Brain Res, 2004, 157:162—173.
- [9] Cirstea MC, Ptiito A, Levin MF. Arm reaching improvements with short-term practice depend on the severity of the motor deficit in stroke[J]. Exp Brain Res, 2003, 152:476—488.
- [10] Cirstea C, Levin M. Improvement of arm movement patterns and endpoint control depends on type of feedback during practice in stroke survivors[J]. Neurorehabil Neural Repair, 2007, 21:398—411.
- [11] 越焕, 单述刚, 谭维溢. 脑卒中后躯干屈曲的动力学分析[J]. 神经损伤与功能重建, 2008, 3(6):441—443.
- [12] Lin KC, Chen YA, Chen CL, et al. The effects of bilateral arm training on motor control and functional performance in chronic stroke: a randomized controlled study[J]. Neurorehabil Neural Repair, 2010,24(1):42—51.
- [13] Caurraugh JH, Lodha N, Naik SK, et al. Bilateral movement training and stroke motor recovery progress: a structured review and meta-analysis[J]. Hum Mov Sci, 2010, 29(5):853—870.
- [14] Stinear CM, Barber PA, Coxon JP, et al. Active-Passive bilateral therapy enhances the effects of upper limb therapy in chronic stroke[J]. Journal of Clinical Neuroscience, 2009,16(3): 465—466.
- [15] 郑雅丹, 胡昔权, 李奎, 陈颖蓓, 解东风. 双侧上肢训练在脑卒中患者康复中的应用[J]. 中国康复医学杂志, 2011, (6):523—528.
- [16] 郑雅丹, 胡昔权. 双侧上肢训练在脑卒中患者康复中应用的研究进展[J]. 中国康复医学杂志, 2011, (3):296—299.
- [17] Messier S, Bourbonnais D, Desrosiers J, et al. Kinematic analysis of upper limbs and trunk movement during bilateral movement after stroke[J]. Arch Phys Med Rehabil, 2006, 87(11):1463—1470.
- [18] 毕胜, 许云影. 偏瘫上肢痉挛、联合反应与异常协同运动临床相关研究[J]. 中国康复医学杂志, 2008, 23(11): 1007—1009.
- [19] 王凯, 陈文锋, 周文益. 脑卒中患者平衡功能损伤的相关因素分析[J]. 中国康复医学杂志, 2007, 22(4):345—346.
- [20] 廖亮华, 江兴妹, 叶志卫, 等. 早期躯干与骨盆控制训练对偏瘫患者运动功能的影响[J]. 中国康复医学杂志, 2011, 26(5):443—446.

