

- years after THA[J]. Acta Orthop, 2010, 81(2):183—188.
- [3] Nantel J, Termoz N, Ganapathi M, et al. Postural balance during quiet standing in patients with total hip arthroplasty with large diameter femoral head and surface replacement arthroplasty[J]. Arch Phys Med Rehabil, 2009, 90(9):1607—1612.
- [4] 吴博,张雷,庞文君,等.老年髋部骨折股骨近端髓内钉—螺旋刀片固定与关节置换术后平衡能力的比较[J].中国康复医学杂志, 2014,29(10):950—952.
- [5] Truszczyńska A, Drzał-Grabiec J, Rapała K, et al. Postural stability in patients with osteoarthritis of the hip. Pilot study [J]. Ortop Traumatol Rehabil, 2013, 15(6):567—573.
- [6] 毕霞,成鹏,吴岳嵩,等.全髋关节置换术后等速测试的临床研究[J].中国康复医学杂志,2005,20(7):520—521.
- [7] 刘玉平,周勇刚,俞光荣.软组织平衡在全髋关节置换术中的重要性[J].中华关节外科杂志(电子版),2009,(6):769—773.
- [8] Sirca A, Susec-Michieli M. Selective type II fibre muscular atrophy in patients with osteoarthritis of the hip[J]. J Neurol Sci, 1980, 44(2—3):149—159.
- [9] Fiori MG, Andreola S, Ladelli G, et al. Selective atrophy of the type IIb muscle fibers in rheumatoid arthritis and progressive systemic sclerosis (scleroderma). A biopsy histochemical study[J]. Eur J Rheumatol Inflamm, 1983, 6(2): 168—181.
- [10] Van Emmerik RE, Remelius JG, Johnson MB, et al. Postural control in women with multiple sclerosis: effects of task, vision and symptomatic fatigue[J]. Gait Posture, 2010, 32(4):608—614.
- [11] Sell TC. An examination, correlation, and comparison of static and dynamic measures of postural stability in healthy, physically active adults[J]. Phys Ther Sport, 2012, 13(2):80—86.
- [12] Missaoui B, Portero P, Bendaya S, et al. Posture and equilibrium in orthopedic and rheumatologic diseases[J]. Neurophysiol Clin, 2008, 38(6):447—457.
- [13] Winiarski S, Czamara A. Evaluation of gait kinematics and symmetry during the first two stages of physiotherapy after anterior cruciate ligament reconstruction[J]. Acta Bioeng Biomech, 2012, 14(2):91—100.
- [14] Błażkiewicz M, Wit A. Comparison of sensitivity coefficients for joint angle trajectory between normal and pathological gait[J]. Acta Bioeng Biomech, 2012, 14(1):83—91.
- [15] Łapszo J, Giovanis V, Prusik K, et al. Balance control contributors - the relationships between leg strength and balance control ability in seniors[J]. Acta Bioeng Biomech, 2012, 14(3):3—8.
- [16] Lenaerts G, Mulier M, Spaepen A, et al. Aberrant pelvis and hip kinematics impair hip loading before and after total hip replacement[J]. Gait Posture, 2009, 30(3):296—302.
- [17] van den Heuvel MR, van Wegen EE, de Goede CJ, et al. The effects of augmented visual feedback during balance training in Parkinson's disease: study design of a randomized clinical trial[J]. BMC Neurol, 2013, (13):137.
- [18] Duncan PW, Weiner DK, Chandler J, et al. Functional reach: a new clinical measure of balance[J]. J Gerontol, 1990, 45(6):M192—197.
- [19] Asayama I, Chamnongkitch S, Simpson KJ, et al. Reconstructed hip joint position and abductor muscle strength after total hip arthroplasty[J]. J Arthroplasty, 2005, 20(4): 414—420.

北京大学第十二届全国肌电图与临床神经生理学习班招生通知

北京大学第三医院神经内科主办的“北京大学第十二届全国肌电图与临床神经生理学习班”拟定于2015年8月10日至14日在北京举办。本届学习班集中了北京大学临床电生理学领域雄厚的师资力量,并邀请国内知名专家共同参与授课。学习班内容包括肌电图、神经传导测定、诱发电位等技术在神经系统疾病诊断治疗中的应用,以及在骨科、手外科、神经外科、运动医学、康复等相关疾病的术中监护与应用。国家级继续医学教育 I 类 8 学分。

Email: bssn1108@sina.com, 联系电话: 010-82264446, 15901312366, 张华纲。

北京大学第三医院神经内科