

参考文献

- [1] DiSipio T, Rye S, Newman B, et al. Incidence of unilateral arm lymphoedema after breast cancer: A systematic review and meta-analysis[J]. The Lancet Oncology, 2013, 14(6):500—515.
- [2] 刘宁飞. 淋巴水肿—诊断与治疗[M].北京:科学出版社,2014. 23—24.
- [3] Morrell RM, Halyard MY, Schild SE, et al. Breast cancer related lymphedema[J]. Mayo Clin Proc,2005,80:1480—1484.
- [4] Kilbreath SL, Refshauge KM, Beith JM, et al. Risk factors for lymphoedema in women with breast cancer: A large prospective cohort[J]. Breast, 2016, 28:29—36.
- [5] Li CZ, Zhang P, Li RW, et al. Axillary lymph node dissection versus sentinel lymph node biopsy alone for early breast cancer with sentinel node metastasis: A meta-analysis[J]. European Journal of Surgical Oncology, 2015, 41(8):958.
- [6] Schmitz KH,Ahmed RL,Troxel AB,et al. Weight lifting for women at risk for breast cancer-related lymphedema: a randomized trial [J]. JAMA,2010,304(24):2699—2705.
- [7] Cheema BS, Kilbreath SL, Fahey PP, et al. Safety and efficacy of progressive resistance training in breast cancer: a systematic review and meta-analysis[J]. Breast Cancer Res Treat,2014,148(2):249—268.
- [8] 陈波,贾实,张文海,等. 乳腺癌术后上肢淋巴水肿的危险因素分析[J]. 中国医科大学学报,2012,28(7):637—641.
- [9] Helyer LK, Varnic M, Le LW, et al. Obesity is a risk factor for developing postoperative lymphedema in breast cancer patients[J]. Breast Journal,2010,16(1):48—54.
- [10] Armer JM, Stewart BR. A comparison of four diagnostic criteria for lymphedema in a post-breast cancer population[J]. Lymphat Res Biol,2005,3(4):208—217.
- [11] Bernas MJ, Askew RL, Armer JM, et al. Lymphedema: How do we diagnose and reduce the risk of this dreaded complication of breast cancer treatment?[J]. Current Breast Cancer Reports,2010,2(1):53—58.
- [12] Ward LC, Czerniec S, Kilbreath SL. Quantitative bioimpedance spectroscopy for the assessment of lymphoedema[J]. Breast Cancer Res Treat,2009,117(3):541—547.
- [13] Cornish B. Bioimpedance analysis: scientific background [J]. Lymphat Res Biol, 2006, 4(1): 47—50.
- [14] Huang TW, Tseng SH, Lin CC, et al. Effects of manual lymphatic drainage on breast cancer-related lymphedema: a systematic review and meta-analysis of randomized controlled trials[J]. World J Surg Oncol,2013;11:15.
- [15] Shao Y, Zhong DS. Manual lymphatic drainage for breast cancer-related lymphoedema[J]. Eur J Cancer Care (Engl), 2017, 26(5):1—5.
- [16] Johansson K, Albertsson M, Ingvar C, et al. Effect of compression bandage with or without manual lymph drainage in patients with postoperative arm lymphedema[J]. Lymphology,1999,32(3):103—110.
- [17] Rucigaj TP, Huljev D. The role of short-stretch bandage in therapy of lymphedema[J]. Acta Med Croatica,2012,66(Suppl 1):49-51.
- [18] Jones LW, Liang Y, Pituskin EN, et al. Effect of exercise training on peak oxygen consumption in patients with cancer: a meta-analysis[J]. Oncologist,2011,16(1):112—120.
- [19] Zeitei C, Yenchun L, Yutzu D. The problems of family caregiving among discharged patients in first month[J]. Nursing Research-Taipei,1999, 7(5):423—434.
- [20] 俞桃英. 护士对出院患者实施电话随访的体会[J]. 中华护理杂志,2006,41(3):246—247.

2018年北京大学第一医院举办全国培训班通知 常见关节肌肉伤病的康复评定和康复治疗技术学习班 ——腰部疾患的循证预防与康复[国家级继续教育项目:2018-16-00-034(国)]

北京大学第一医院康复医学科将在2018年6月20—23日举办第11届常见关节肌肉伤病的康复评定和康复治疗技术学习班。本届学习班将在循证医学基础上结合现代化康复手段,以全新的腰痛康复理念和实用的康复技术为学员呈现一场腰痛康复盛宴。学习班将注重临床理论与实践相结合,强调实用性。学习班特别邀请最新专著《腰部疾患:循证预防与康复》中文版的三位主编,北京大学第一医院王宁华主任、北京医院顾新主任和中日友好医院谢欲晓主任携主要译者亲自授课;还特别邀请曾任台湾仁爱医院院长、台湾神经肌肉整合系统康复技术规范化疗体系创始人的黄文成教授来京授课,将悬吊技术整合在课程中。

学习班学费3000元(包含最新译著《腰部疾患:循证预防与康复》1本)。提供午餐,其他食宿、交通费用自理。5月31日前报名可享受早鸟价2600元。考核合格者授予国家级继续教育I类学分6分。限额50人,先报先得。

报名方式:微信扫描以下二维码直接在线报名。

咨询电话:010-83572455或010-83575162。



报名二维码